### **OHIC Aligned Primary Care Measure Set**

Effective Date: Performance Period Starting on or after 1-1-21

Total Measures (excluding Developmental): 21 Total Measures (including Developmental): 28

### Core Measures by Domain (7)

### Chronic Illness (2)

- Comprehensive Diabetes Care: Eye Exam
- Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Control (<8.0%)

## Preventive Care (5)

- Breast Cancer Screening
- Colorectal Cancer Screening
- Developmental Screening in the First Three Years of Life
- Lead Screening for Children
- Weight Assessment and Counseling for Nutrition and Physical Activity for Children/ Adolescents

# Menu Measures by Domain (14)

## Behavioral Health (2)

- Depression Remission at Six Months
- Unhealthy Drug and Alcohol Use Screening and Brief Counseling

# Chronic Illness (2)

- Controlling High Blood Pressure
- Comprehensive Diabetes Care: Medical Attention for Nephropathy
- Kidney Health Evaluation for Patients with Kidney Disease

## Consumer Experience (1)

• PCMH CAHPS (for primary care) - questions not specified

## Preventive Care (6)

- Adolescent Well-Care Visits
- Adult BMI Assessment
- Cervical Cancer Screening
- Child and Adolescent Well-Care Visits (Adolescent Well-Care Visits only)
- Childhood Immunization Status (Combo 10)
- Chlamydia Screening
- Fluoride Varnish
- Immunizations for Adolescents (Combo 2)

# Maternity Care (1)

• Maternal Depression Screening

## Utilization (2)

- Emergency Department Visits per 1,000
- Inpatient Visits per 1,000 (Inpatient Utilization General Hospital/Acute Care)

# **Developmental Measures by Domain (7)**

## Behavioral Health (3)

- Concurrent Use of Opioids and Benzodiazepines
- Depression Remission or Response for Adolescents and Adults
- Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults

# Chronic Illness (1)

• Statin Therapy for Patients with Cardiovascular Disease

## Preventive Care (2)

- Depression Screening and Follow-Up for Adolescents and Adults
- Unhealthy Alcohol Use Screening and Follow-Up

# Social Determinants of Health (1)

### **OHIC Aligned Maternity Care Measure Set**

Effective Date: Performance Period Starting on or after 1-1-21 Total Measures (excluding Developmental): 7 Total Measures (including Developmental): 7

# Core Measures by Domain (3)

### Prenatal Care (2)

- Behavioral Health Risk Assessment Screenings
- Prenatal and Postpartum Care: Timeliness of Prenatal Care

# Postpartum Care (1)

• Prenatal and Postpartum Care: Postpartum Care

# Menu Measures by Domain (4)

# Labor and Delivery (2)

- Cesarean Rate for Nulliparous Singleton Vertex (PC-02)
- Elective Delivery Prior to 39 Completed Weeks Gestation (PC-01)

# Postpartum Care (1)

• Maternity Care: Post-Partum Follow-Up and Care Coordination

## Neonatal Care - Term Infants (1)

• Exclusive Breast Milk Feeding (PC-05)

### OHIC Aligned Outpatient Behavioral Health - Mental Health Measure Set

Effective Date: Performance Period Starting on or after 1-1-21

Total Measures (excluding Developmental): 10 Total Measures (including Developmental): 14

### Core Measures by Domain (0)

## Menu Measures by Domain (10)

### Acute Care (2)

- Adult Major Depressive Disorder (MDD): Coordination of Care of Patients with Specific Comorbid Conditions
- Depression Remission at Six Months

### Chronic Illness Care (2)

- Diabetes Screening for People with Schizophrenia or Bipolar Disorder who Are Using Antipsychotic Medications
- Metabolic Monitoring for Children and Adolescents on Antipsychotics

## Hospital (2)

- Follow-Up After Emergency Department Visit for Mental Illness
- Follow-Up After Hospitalization for Mental Illness (7-Day) (NCQA Version)

# Medication Management (1)

• Antidepressant Medication Management

### Prevention (3)

- Adult Major Depressive Disorder (MDD): Suicide Risk Assessment
- Child and Adolescent Major Depressive Disorder: Suicide Risk Assessment
- Unhealthy Drug and Alcohol Use: Screening & Brief Counseling

## **Developmental Measures by Domain (4)**

### Acute Care (2)

- Depression Remission or Response for Adolescents and Adults
- Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults

# Preventive Care (1)

Unhealthy Alcohol Use Screening and Follow-Up

### Social Determinants of Health (1)

# OHIC Aligned Outpatient Behavioral Health - Substance Use Treatment Measure Set

Effective Date: Performance Period Starting on or after 1-1-21

Total Measures (excluding Developmental): 2 Total Measures (including Developmental): 4

## Core Measures by Domain (1)

### Chronic Care Illness (1)

• Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment

## Menu Measures by Domain (1)

## Hospital (1)

• Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence

# **Developmental Measures by Domain (2)**

### Behavioral Health (1)

• Concurrent Use of Opioids and Benzodiazepines

## Social Determinants of Health (1)

### **OHIC Aligned Acute Care Hospital Measure Set**

Tentative Effective Date: Performance Period Starting on or after 1-1-21 Total Measures (excluding Developmental): 16 Total Measures (including Developmental): 17

## Core Measures by Domain (8)

### Hospital (8)

- CAUTI: Catheter-Associated Urinary Tract Infection (HAI-2)
- CLABSI: Central Line-Associated Blood Stream Infection (HAI-1)
- Clostridium Difficile (C.diff.) Infections (HAI-6)
- Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence
- Follow-Up After Emergency Department Visit for Mental Illness
- Follow-Up After Hospitalization for Mental Illness (7-Day) (NCQA Version)
- HCAHPS
- Hospital-wide Readmit (READM-30-HOSP-WIDE)

### Menu Measures by Domain (8)

# Hospital (7)

- Alcohol & Other Drug Use Disorder Treatment Provided or Offered at Discharge (SUB-3) and Alcohol & Other Drug Use Disorder Treatment at Discharge (SUB-3a)
- American College of Surgeons Centers for Disease Control and Prevention (ACS-CDC)
   Harmonized Procedure Specific Surgical Site Infection (SSI) Outcome Measure HAI-3:

   SSI: Colon Surgical Site Infection for Colon Surgery and HAI-4: SSI: Hysterectomy Surgical Site Infection for Abdominal Hysterectomy
- Cesarean Rate for Nulliparous Singleton Vertex (PC-02)
- Elective Delivery Prior to 39 Completed Weeks Gestation (PC-01)
- Exclusive Breast Milk Feeding (PC-05)
- Methicillin-resistant Staphylococcus Aureus (or MRSA) Blood Infections (HAI-5)
- Severe Sepsis and Septic Shock: Management Bundle

# Population Health (1)

• 30-Day All-Cause Unplanned Readmission Following Psychiatric Hospitalization (CMS Version)

## **Developmental Measures by Domain (1)**

### Social Determinants of Health (1)

### OHIC Aligned Behavioral Health Hospital Measure Set

Tentative Effective Date: Performance Period Starting on or after 1-1-21 Total Measures (excluding Developmental): 9 Total Measures (including Developmental): 9

# Core Measures by Domain (2)

# Hospital (2)

- Follow-Up After Hospitalization for Mental Illness (7-Day) (NCQA Version)
- Timely Transmission of Transition Record

## Menu Measures by Domain (7)

### Hospital (6)

- Alcohol & Other Drug Use Disorder Treatment Provided or Offered at Discharge (SUB-3) and Alcohol & Other Drug Use Disorder Treatment at Discharge (SUB-3a)
- Hours of Physical Restraint Use (HBIPS-2)
- Hours of Seclusion Use (HBIPS-3)
- Patients Discharged on Multiple Antipsychotic Medications with Appropriate Justification (HBIPS-5)
- Screening for Metabolic Disorders
- Transition Record with Specified Elements Received by Discharged Patients

## Population Health (1)

• 30-Day All-Cause Unplanned Readmission Following Psychiatric Hospitalization (CMS Version)

### **OHIC Aligned ACO Measure Set**

Effective Date: Performance Period Starting on or after 1-1-21 Total Measures (excluding Developmental): 36

Total Measures (including Developmental): 43

## Core Measures by Domain (7)

## Chronic Care Illness (2)

- Comprehensive Diabetes Care: Eye Exam
- Comprehensive Diabetes Care: HbA1c Control (<8.0%)

### Hospital (1)

• Follow-Up After Hospitalization for Mental Illness (7-Day) (NCQA Version)

### Preventive Care (4)

- Breast Cancer Screening
- Colorectal Cancer Screening
- Developmental Screening in the First Three Years of Life
- Weight Assessment and Counseling for Nutrition and Physical Activity for Children/ Adolescents

## Menu Measures by Domain (29)

# Behavioral Health (1)

Depression Remission at Six Months

### Care Coordination (1)

• Advance Care Plan (CMS Version)

### Chronic Illness (3)

- Comprehensive Diabetes Care: Medical Attention for Nephropathy
- Controlling High Blood Pressure
- Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment
- Kidney Health Evaluation for Patients with Kidney Disease

# Hospital (10)

- Cesarean Rate for Nulliparous Singleton Vertex (PC-02)
- Elective Delivery Prior to 39 Completed Weeks Gestation (PC-01)
- Exclusive Breast Milk Feeding (PC-05)
- Follow-Up After Emergency Department Visit for Alcohol and Other Drug Dependence
- Follow-Up After Emergency Department Visit for Mental Illness
- HCAHPS
- Hospital-wide Readmit (READM-30-HOSP-WIDE)

- Plan (ACO) All-Cause Readmission
- Timely Transmission of Transition Record
- Transition Record with Specified Elements Received by Discharged Patients

### Overuse (1)

• Use of Imaging Studies for Low Back Pain

# Preventive Care (13)

- Adolescent Well-Care Visit
- Adult BMI Assessment
- Adult Major Depressive Disorder (MDD): Suicide Risk Assessment
- Behavioral Health Risk Assessment Screenings
- Cervical Cancer Screening
- Child and Adolescent Major Depressive Disorder: Suicide Risk Assessment
- Child and Adolescent Well-Care Visits (Adolescent Well-Care Visits only)
- Childhood Immunization Status (Combo 10)
- Chlamydia Screening
- Fluoride Varnish
- Immunizations for Adolescents (Combo 2)
- Lead Screening in Children
- Maternal Depression Screening
- Prenatal & Postpartum Care Postpartum Care Rate
- Unhealthy Drug and Alcohol Use: Screening & Brief Counseling

## Developmental Measures by Domain (7)

### Behavioral Health (3)

- Concurrent Use of Opioids and Benzodiazepines
- Depression Remission or Response for Adolescents and Adults
- Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults

### Chronic Illness (1)

• Statin Therapy for Patients with Cardiovascular Disease

# Preventive Care (2)

- Depression Screening and Follow-Up for Adolescents and Adults
- Unhealthy Alcohol Use Screening and Follow-Up

### Social Determinants of Health (1)